

Case Report

Usage of Trapezoid RX Wire guided Retrieval Basket with the Alliance™ II Inflation Handle for Treatment of Bouveret Syndrome: A Rare Clinical Entity.

Mahmod Ghoname ¹, Hazem Abosheishaa ^{2*}, Arshia Sethi ², Sharif Abbas ³, Sherif Hegab ⁴, Moataz Yousry Soliman ¹

1- Hepatology, Gastroenterology and Infectious Diseases Department, Al-Azhar University, Faculty of Medicine for Boys, Cairo, Egypt.

2- Internal Medicine Department, Icahn School of Medicine at Mount Sinai, NYC Health + Hospitals Queens, NY, USA.

3- Internal Medicine Department, Koc University School of Medicine, Istanbul, Turkey.

4- Hepatology, Gastroenterology and Infectious Diseases Department, Ahmed Maher Teaching Hospital, GOTHI, Cairo, Egypt.

ARTICLE INFO

Article history:

Received 14 Dec. 2024

Received in revised form 24 Dec. 2024

Accepted 27 Dec. 2024

Keywords:

Gallstones

Gastric outlet obstruction

Bouveret syndrome

Gastrointestinal

ABSTRACT

Bouveret Syndrome is an infrequent manifestation of gallstone disease causing gastric outlet obstruction. This syndrome has diagnostic and therapeutic challenges due to its rarity and non-specific clinical presentation. Here, we present a case report involving a 78-year-old female patient who experienced gastric outlet blockage due to the presence of a sizable gallstone. A novel technique was applied wherein a trapezoid RX wire-guided retrieval basket (Boston Scientific) was utilized with Esophagogastroduodenoscopy to remove the stone without the need for surgical intervention. Post-procedure monitoring demonstrated a smooth recovery with immediate relief of symptoms. Imaging confirmed the absence of residual gallstones or obstruction.

* **Corresponding author:** Hazem Abosheishaa, Department of Medicine, Icahn School of Medicine at Mount Sinai, NYC Health + Hospitals Queens, NY, USA.. Email Address: <mailto:hazemabosheishaa@gmail.com>

ISSN (Print) 3066-4004, ISSN (Online) 3066-4012– see front matter © 2024 ASIDE Gastroenterology. This work is licensed under a Creative Commons Attribution 4.0 International License. Hosting by ASIDE Journals. All rights reserved.

Citation: Ghoname M, Abosheishaa H, Sethi A, Abbas S, Hegab S, Soliman MY. Usage of Trapezoid RX Wire guided Retrieval Basket with the Alliance™ II Inflation Handle for Treatment of Bouveret Syndrome: A Rare Clinical Entity. ASIDE GI. 2025;1(1):1-4. doi:10.71079/ASIDE.GI.0102255

<https://doi.org/10.71079/ASIDE.GI.0102255>

Journal homepage: <https://asidejournals.com/index.php/Gastroenterology>

1. Introduction

Bouveret syndrome (BS) is an uncommon manifestation of gallstone disease characterized by gastric outlet obstruction where the gallstone gets lodged in the duodenum or pylorus [1]. BS has been linked to a high mortality rate of up to 27% and can cause serious complications, particularly in elderly patients or those with comorbidities. Due to the rarity and non-specific presentation of BS, diagnosing it can be difficult [2]. A correct and prompt diagnosis is essential for enhancing prognosis, lowering morbidity, and preventing death [3]. Here, we present a case of BS with a large gallstone causing gastric outlet obstruction for which a trapezoid RX wire-guided retrieval basket (Boston Scientific) is used during EGD without surgical intervention.




Figure 1: Multislice CT of the abdomen with oral and IV contrast showing partially distended irregular gall bladder containing multiple gas vacuoles (green arrows) which are seen compressing the second part of the duodenum (black arrows) with a fistulous track in between exerting proximal dilation of the stomach. Obvious pneumobilia is also seen.

On initial laboratory evaluation, she had a hemoglobin level of 9.3 g/dL and a white cell count of $15.3 \times 10^9/L$ with elevated CRP 12 mg/dl. Serum creatinine was 1.4 mg/dL, and random blood sugar was 295 mg/dL with no acetone in urine. Her amylase and lipase were 52 U/L and 65 U/L, respectively. Bilirubin was normal with an alkaline phosphatase (ALP) level of 155 U/l (normal range, 40–120 U/l) and a γ -glutamyl transferase (GGT) level of 99 U/l (normal, <37 U/l). Diagnostic workup, including an abdominal X-ray and CT scan, confirmed the presence of a large gallstone causing gastric outlet obstruction with a fistulous tract between the gallbladder and the duodenum and a dilated proximal stomach (Figure 1).

After careful assessment and evaluation, an upper GI endoscopy was performed to relieve the gastric outlet obstruction. The gastroscope was inserted after endotracheal intubation, and the obstructing gallstone was visualized in the pyloric ring obstructing the lumen (Figure 2). We decided to use the trapezoid RX wire-guided retrieval basket (Boston Scientific), designed for crushing and removing stones in the biliary duct. We may use the Alliance™ II Inflation Handle for mechanical lithotripsy to crush large stones trapped within the basket with the scope in place. However, after several trials, stone extraction failed. So, we decided to try to mobilize the large stone to the stomach. The biliary extractor balloon catheter was inflated up to 18 mm distal to obstructing stone, and the stone was successfully retrieved into the stomach with difficulty. A trapezoid basket was introduced through the endoscope's

2. Case Report

A 78-year-old female patient presented with a two-month history of worsening abdominal pain, nausea, and vomiting. The patient reported a decreased appetite and unintentional weight loss but no history of hematemesis, melena, dysphagia, fever, or rigors. Her medical history included type 2 diabetes mellitus, hypertension, ischemic heart disease, atrial fibrillation, NSAID due to sciatica, and a history of chronic calculous cholecystitis with an unremarkable surgical history. She was afebrile, slightly hypertensive, and with an irregular heart rate. Physical examination revealed abdominal distension and tenderness in the epigastric region with no organomegaly.

working channel and used to crush, grasp, and extract the large gallstone, which measures 4 cm \times 3.6 cm, and to facilitate the extraction of the stone through the cardia and esophagus without impaction or perforation. The remaining stones were retrieved without any complications (Figure 3). Finally, gastric outflow was restored after three hours of working in a single session.

The patient experienced immediate relief and post-procedure monitoring showed a steady recovery without any complications. The patient tolerated a regular diet, and subsequent imaging studies confirmed the absence of residual gallstones or signs of obstruction.

3. Discussion

Bouveret's syndrome (BS) is a rare subtype of gallstone ileus making up about 1%-3% of cases [4]. Repeated episodes of cholecystitis result in inflammation and adhesions between the gallbladder and the GI tract which most commonly involves the duodenum but can also involve the stomach. Pressure necrosis by fairly large calculus results in fistula formation which predisposes to a migrating calculus hence causing either obstruction or ileus [5].

BS is usually seen in elderly females, with a median age of 74 and a history of cholelithiasis or cholecystitis. Patients present with vague symptoms of gastric outlet or bowel obstruction that usually include nausea or vomiting and sometimes hematemesis or melena [5]. Non-

specific presenting symptoms not only make diagnosis challenging but also increase morbidity and mortality [4].




Figure 2: Endoscopic picture showing a large stone obstructing the pyloric ring with a trial to remove it using a Retrieval Basket (Boston Scientific).




Figure 3: Showing the pyloric ring, duodenal bulb, and second part of the duodenum after the removal of the stone.

Multiple diagnostic modalities can be used in the workup for BS, but CT and EGD are the most widely utilized diagnostic modalities with the highest specificity and sensitivity, (95% and 100% in the former) with EGD being the most commonly used modality in patients presenting with overt bleeding. CT with contrast can also be used in diagnosis with the extravasated contrast outlining the fistula [6].

In our case, Abdominal X-rays followed by CT were sufficient to reach a definitive diagnosis of BS.

In terms of BS management, endoscopy utilization is especially worthwhile in elderly patients with significant comorbidities [7]. However, a study done by Howells et al. showed up to 91% failure of endoscopic and percutaneous extraction despite the availability of expert providers [8]. Nevertheless, a variety of different endoscopic techniques have been shown to be successful [9]. For example, Endoscopic Electrohydraulic Lithotripsy (EHL) has been utilized by Avci et al. to help dissolve a 3 cm biliary stone lodged at the proximal duodenum [10]. In another case, gastric outflow was restored endoscopically in a poor surgical candidate with a pigmented gallstone using Roth Net Platinum Universal Retriever [11].

In case EGD fails, surgery is required. Traditional methods like open gastrotomy, pylorotomy, or duodenotomy are often used but are less preferred due to higher morbidity and mortality rates. Laparoscopy is also used in some cases where open surgery is not an option. For elderly patients who are not suitable candidates for surgery due to increased surgical risks, a two-step procedure is considered. This involves first removing the stone and then performing cholecystectomy and fistula repair [12].

4. Conclusions

In conclusion, using the trapezoid RX wire-guided retrieval basket (Boston Scientific) along with Alliance™ II Inflation Handle for mechanical lithotripsy and the biliary extractor balloon catheter as described in this endoscopic approach to treat BS may improve patient outcomes.

Conflicts of Interest:

None

Ethical Approval:

None

Funding Source:

None

Acknowledgements:

None

Institutional Review Board (IRB) Approval:

None

LLM Statement:

None

Authors Contribution Statement:

MG wrote the case presentation section, case reporter, and provided the endoscopy images. HA reviewed the previous literature, wrote the introduction, extracted some key points, contributed to the discussion writing and corresponding author. AS contributed to the discussion section by describing previous case data. SA Wrote the main discussion part after reviewing the literature, and submission of the article. SH reviewed the manuscript and wrote the abstract. MS Whole paper review, senior author of the paper.

Data Availability Statement:

Full Endoscopic video of the case is available with the author upon request.

References

1. Turner AR, Kudaravalli P, Al-Musawi JH, Ahmad H. Bouveret Syndrome (Bilioduodenal Fistula). StatPearls. Treasure Island (FL): StatPearls Publishing, Copyright © 2023, StatPearls Publishing LLC.; 2024.
2. Dunlop H, Goodarzi MR. Bouveret Syndrome: A Rare Case of Gallstone Ileus Further Complicated by Stone Migration. Cureus. 2020; e11219 [PMID: 33154859 10.7759/cureus.11219: 10.7759/cureus.11219]
3. Qasaimeh GR, Bakkar S, Jadallah K. Bouveret's Syndrome: An Overlooked Diagnosis. A Case Report and Review of Literature. Int Surg. 2014; 819 [PMID: 25437593 10.9738/INTSURG-D-14-00087.1: 10.9738/INTSURG-D-14-00087.1]
4. Probert S, Cai W, Islam F, Ballanamada Appaiah NN, Salih A. Bouveret Syndrome: A Rare Case and Review of the Literature. Cureus. 2022; e24768 [PMID: 35686250 10.7759/cureus.24768: 10.7759/cureus.24768]
5. Yu YB, Song Y, Xu JB, Qi FZ. Bouveret's syndrome: A rare presentation of gastric outlet obstruction. Exp Ther Med. 2019; 1813 [PMID: 30783453 10.3892/etm.2019.7150: 10.3892/etm.2019.7150]
6. Nickel F, Muller-Eschner MM, Chu J, von Tengg-Kobligk H, Muller-Stich BP. Bouveret's syndrome: presentation of two cases with review of the literature and development of a surgical treatment strategy. BMC Surg. 2013; 33 [PMID: 24006869 10.1186/1471-2482-13-33: 10.1186/1471-2482-13-33]
7. Zhao JC, Barrera E, Salabat M, Denham W, Leung D, Ujiki M. Endoscopic treatment for Bouveret syndrome. Surg Endosc. 2013; 655

[PMID: 23052513 10.1007/s00464-012-2533-8: 10.1007/s00464-012-2533-8]

8. L H, L L, M D. Gallstone Coleus: A Rare Relation of Gallstone Ileus. International Journal of Surgery and Research. 2015; 28 10.19070/2379-156x-150006: 10.19070/2379-156x-150006]

9. Puri V, Lee RW, Amirlak BA, Lanspa SJ, Fitzgibbons RJ, Jr. Bouveret syndrome and gallstone ileus. Surg Laparosc Endosc Percutan Tech. 2007; 328 [PMID: 17710061 10.1097/SLE.0b013e31806c7dc2: 10.1097/SLE.0b013e31806c7dc2]

10. Avci A, Gogenur I, Bulut M. Bouveret's syndrome treated with endoscopic electrohydraulic lithotripsy. BMJ Case Rep. 2019; [PMID: 30798279 10.1136/bcr-2018-228316: 10.1136/bcr-2018-228316]

11. Jindal A, Philips CA, Jamwal K, Sarin SK. Use of a Roth Net Platinum Universal Retriever for the endoscopic management of a large symptomatic gallstone causing Bouveret's syndrome. Endoscopy. 2016; E308 [PMID: 27669536 10.1055/s-0042-116430: 10.1055/s-0042-116430]

12. Rey Chaves CE, Villamil CJ, Ruiz S, Galvis V, Conde D, Sabogal Olarte JC. Cholecystogastric fistula in Bouveret syndrome: Case report and literature review. Int J Surg Case Rep. 2022; 106918 [PMID: 35339038 10.1016/j.ijscr.2022.106918: 10.1016/j.ijscr.2022.106918]