



Original Article

Improving Inpatient Care for Acute Chronic Obstructive Pulmonary Disease Exacerbations: A Quality Improvement Initiative

Khaqan Ahmed¹, Afaq Ahmad¹, Muhammad Saqib¹, Muhammad Iftikhar², Humna Shahid³

1-Department of Medicine, Khyber Teaching Hospital, Peshawar, Pakistan

2-Department of Medicine, Hayatabad Medical Complex, Peshawar, Pakistan

3-Department of Medicine, CMH Multan Institute of Medical Sciences Multan, Pakistan

ARTICLE INFO

Article history:

Received 13 Jul. 2025

Received in revised form 15 Aug. 2025

Accepted 24 Aug. 2025

Published 20 Sep. 2025

Keywords:

COPD

Acute Exacerbation

Quality Improvement

Electronic Health Record

Pulmonary Rehabilitation

Smoking Cessation

ABSTRACT

Background: Acute exacerbations of chronic obstructive pulmonary disease (AECOPD) account for high hospital use; guideline-concordant diagnostics, oxygen titration, and discharge planning are often under-delivered. We aim to measure baseline compliance with eight AECOPD process standards and evaluate whether EHR-embedded prompts plus staff education were associated with observed improvements.

Methods: We audited 50 consecutive adult admissions with primary AECOPD (ICD-10 J44.1 or clinician-documented diagnosis) between July and October 2024, implemented EHR prompts, an orderset, and targeted education (Nov–Dec 2024), then re-audited 50 consecutive admissions (Jan–Mar 2025) using identical methods. Primary process measures were pre-specified; analyses were descriptive with 95% Wilson confidence intervals (CIs) for proportions and two-sided Fisher exact tests for pre/post comparisons. This work was performed as a service evaluation/pilot.

Results: After implementation, six of eight process standards showed observed increases. Pulmonary rehabilitation referrals rose from 32/50 (64.0%, 95% CI 50.1–75.9%) to 43/50 (86.0%, 95% CI 73.8–93.0%), $p = 0.0198$. Oxygen titration improved from 40/50 (80.0%, 95% CI 66.0–88.6%) to 46/50 (92.0%, 95% CI 78.6–95.7%), $p = 0.148$. Other measures (ABG timing, bronchodilator and corticosteroid delivery, appropriate antibiotic use, CXR timing, smoking advice) showed observed improvements; p -values and CIs are reported in the text.

Conclusions: Embedding guideline-aligned prompts and an orderset in the EHR, together with staff education, was associated with observed improvements in several AECOPD process measures in this pilot QI project. Findings should be interpreted cautiously, given the pilot design, multiple comparisons, and limited sample size; further controlled and longitudinal evaluation is indicated.

1. Introduction

1.1. Problem Description

Chronic obstructive pulmonary disease (COPD) is among the most common chronic illnesses encountered in internal medicine practice, responsible for high rates of morbidity, mortality, and healthcare utilization. Acute exacerbations of COPD (AECOPD) contribute disproportionately to this burden, representing up to 70% of hospital admissions among COPD patients and often resulting in prolonged hospital stays and avoidable readmissions [1, 2, 3, 4]. Within our institution, informal staff discussions and case reviews identified inconsistencies in the inpatient management of AECOPD, particularly in the timely administration of key

therapies, adherence to oxygen titration guidelines, and discharge planning.

1.2. Available Knowledge

National and international guidelines, particularly the GOLD and NICE NG115, outline best practices for AECOPD management [5, 6, 7, 8]. These include early arterial blood gas (ABG) testing, rapid administration of short-acting bronchodilators and corticosteroids, judicious antibiotic use based on sputum characteristics and clinical findings, controlled oxygen delivery targeting SpO_2 88–92%, timely imaging to exclude other diagnoses, and referral to pulmonary rehabilitation. Despite this guidance, studies suggest that real-world practice often diverges from these standards due to a variety of system- and provider-level barriers. Recent reviews and audits have highlighted frequent issues with oxygen overuse, poor documentation, and under-referral to pulmonary rehab—findings consistent with our internal observations [9, 10, 11, 12, 13, 14].

1.3. Rationale

Our team hypothesized that many of the observed care gaps stemmed from workflow inefficiencies and missed clinical prompts during high-pressure admissions. Given that our hospital had already adopted electronic prescribing and documentation systems, we reasoned that integrating clinical decision support tools, such as automatic prompts or checklists, into existing digital workflows

*Corresponding author: Muhamad Saqib, Department of Medicine, Khyber Teaching Hospital, Peshawar, Pakistan. Email: muhammadsaqib.drkmc@gmail.com

Published by the American Society for Inclusion, Diversity, and Equity in Healthcare (ASIDE). ISSN (Print): 3066-4004, ISSN (Online): 3066-4012. © 2025 The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY 4.0). Hosting by ASIDE Journals.

Citation: Ahmed K, Ahmad A, Saqib M, Iftikhar M, Shahid H. Improving Inpatient Care for Acute Chronic Obstructive Pulmonary Disease Exacerbations: A Quality Improvement Initiative. ASIDE Int Med. 2025;2(3):1-10, doi:10.71079/ASIDE.IM.092025173




Figure 1: Quality Improvement Process Flowchart for AECOPD Inpatient Care This flowchart summarizes the sequential steps undertaken in the quality improvement (QI) project to enhance adherence to guideline-based inpatient care for acute exacerbations of chronic obstructive pulmonary disease (AECOPD). Starting with a baseline audit of patient records from July to October 2024, the team identified key process gaps in oxygen titration, pulmonary rehabilitation referral, and imaging timelines. Multidisciplinary interventions, including electronic health record (EHR) prompts, automated discharge tools, and staff education, were implemented between November and December 2024. A re-audit from January to March 2025 was used to evaluate the impact of these changes, informing plans for sustainability and scale-up.

could enhance compliance with evidence-based standards. In addition, educational reinforcement through departmental meetings and targeted communication was anticipated to improve clinical staff engagement and awareness.

1.4. Specific Aims

The primary aim of this quality improvement project was to assess our baseline compliance with key process standards for AECOPD management and implement low-cost, replicable interventions to address identified care gaps. We also aimed to measure the effectiveness of these interventions through a structured re-audit and generate actionable insights for future system-wide changes.

2. Methods

2.1. Organizational context

The project took place on a 650-bed tertiary academic hospital general medicine ward with an existing integrated EHR and in-house clinical informatics support. The clinical team included on-call respiratory specialists, ward physicians, nursing staff, physiotherapy (pulmonary rehabilitation), pharmacy, and an antimicrobial stewardship pharmacist who participated in the QI steering group.

2.2. Patient selection and case identification

We included consecutive adult patients (18 years) admitted to the general medicine ward with a primary diagnosis of acute exacerbation of COPD (AECOPD). Case identification relied on either




Figure 2: Compliance with AECOPD Care Standards Before and After Intervention Bar chart comparing baseline and re-audit compliance rates across eight predefined process standards for inpatient management of acute exacerbations of COPD (AECOPD). Post-intervention gains were observed in all domains, with the statistically significant improvements noted in pulmonary rehabilitation referral and oxygen titration documentation. Standards were benchmarked against national and international guideline targets (NICE, GOLD, BTS, USPSTF). Each bar represents the percentage of patients meeting the criterion during the respective audit cycle (n = 50 per cycle).




Figure 3: Proportion of Standards Meeting Target Compliance Thresholds Post-Intervention Pie chart depicting the proportion of care standards that met or did not meet target benchmarks during the re-audit period (January–March 2025). Six of eight standards achieved full compliance, indicating successful implementation of most intervention strategies. Oxygen titration and chest X-ray turnaround time, though improved, remained below predefined targets, warranting further optimization.

clinician documentation of “COPD exacerbation” in admission or ED records or ICD-10 coding consistent with AECOPD (J44.1). Exclusion criteria were: primary diagnosis other than AECOPD, invasive mechanical ventilation instituted prior to the EHR trigger or admission for palliative care only. Some process measures had reduced denominators due to documentation availability (for example, antibiotic indication was not appraisable in 3 baseline patients, and smoking status was unavailable in 10 patients); we report the denominators for each measure in Tables 2–3 and describe the handling of incomplete documentation in the Results.

2.3. Interventions

Based on root cause analysis and staff feedback, four main interventions were introduced between November and December 2024. First, we implemented an oxygen prescription checklist embedded into the electronic prescribing system to improve documentation of target SpO₂ ranges and FiO₂. Second, we developed an automatic referral prompt for pulmonary rehabilitation that triggered at discharge for eligible patients. Third, we delivered guideline-focused education sessions to all medical and nursing staff. Finally, a pop-up reminder was added to the discharge summary module to




Figure 4: Action Plan Implementation Progress: Baseline, Target, and Re-Audit Comparison Grouped bar chart showing baseline status, target compliance thresholds, and re-audit performance for four key intervention metrics: oxygen titration compliance, staff education coverage, pulmonary rehabilitation referral, and smoking cessation documentation. Staff education, pulmonary rehab referrals, and smoking cessation advice achieved or exceeded targets. Oxygen titration compliance improved but remained marginally below the 95% target.

ensure that documentation of smoking cessation advice is provided for current smokers. These interventions were co-designed with the respiratory team, clinical informatics unit, and physiotherapy department.

2.4. Acute inhaled therapy

The electronic orderset recommended a metered-dose inhaler (MDI) with a spacer as the preferred bronchodilator delivery method because of equivalent efficacy and logistical advantages (including lower aerosol generation). Nebulized bronchodilator therapy remained available and was permitted when MDI/spacer was not feasible or tolerated (for example, where patients were unable to coordinate inhalation, were highly agitated, or at the clinician's discretion). We clarified this preference within the orderset and included a Table 1 footnote to reflect the local policy.

2.5. Antibiotic decision-making and stewardship.

The EHR prompt did not automatically prescribe antibiotics. Instead, the prompt and orderset presented guideline-based criteria to prompt clinician consideration of antibiotics (presence of increased sputum purulence, increased sputum volume, increased dyspnoea, temperature $> 38^{\circ}\text{C}$, or radiographic consolidation). Final prescribing decisions remained at the discretion of the treating clinician. The hospital antimicrobial stewardship pharmacist subsequently reviewed all antibiotic prescriptions for the audit cohorts during the QI period, and stewardship feedback was communicated to treating teams when prescriptions appeared discordant with the indicated criteria.

2.6. Systemic corticosteroids.

The orderset included a recommended short-course systemic corticosteroid regimen consistent with contemporary guideline practice: oral prednisolone 40 mg once daily for 5 days (or an equivalent intravenous regimen when oral administration was not possible). An informational notice within the orderset prompted clinicians to review recent corticosteroid prescriptions in the record to avoid unnecessary cumulative oral corticosteroid exposure and to discourage prolonged tapers unless clinically justified.

2.7. Implementation and target selection.

Targets for each process indicator were selected by a multidisciplinary QI steering group (respiratory physicians, ward physicians, nurses, pharmacy, physiotherapy, and clinical informatics) using a pragmatic balance of ambition and feasibility informed by baseline performance. We used staged compliance thresholds (85% $\rightarrow$ 90% $\rightarrow$ 95%) to secure early, achievable wins and then progressively raise them in subsequent Plan-Do-Study-Act (PDSA) cycles. The choice of these operational targets was therefore locally informed rather than derived from a single external numerical standard; this approach followed the Model for Improvement and aimed to sustain momentum through iterative improvement.

2.8. QI framework and iterative approach.

The QI project employed the Model for Improvement, utilizing rapid Plan-Do-Study-Act (PDSA) cycles. A multidisciplinary team (physicians, nurses, pharmacists, physiotherapists, and informatics specialists) met weekly during the implementation to review process measures, frontline feedback, and iterate on prompt wording, order defaults, and workflow. Figure 1 summarises the PDSA timeline.

2.9. Study of the Interventions

We employed a before-and-after audit design to assess the effectiveness of the intervention. Data were collected retrospectively for the baseline cohort (July–October 2024) and prospectively for the re-audit cohort (January–March 2025). Observed outcomes were compared to established targets to assess changes in compliance. Attribution of outcomes to interventions was supported by temporality, stability of external conditions, and qualitative feedback from clinical staff. Allocation and blinding. This was an uncontrolled before-and-after QI audit, and there was no randomization or blinding. Staff were aware of the interventions and audits (education sessions and EHR prompts), which may have influenced behaviour (see Limitations). To enhance transparency and reproducibility, a visual summary of the QI cycle is provided in Figure 1. This flowchart outlines the sequential steps of the quality improvement process, beginning with the baseline audit conducted

Table 1: Audit standards and targets

No.	Standard	Source	Target
1	ABG within 1 hour	NICE NG115 [7]	≥90%
2	Nebulized bronchodilators within 1 hour	GOLD [8]	≥90%
3	Corticosteroids within 4 hours	GOLD; NICE [5, 7, 8]	≥95%
4	Antibiotics only if indicated	NICE NG115 [7]	≥90%
5	Oxygen titration to SpO ₂ 88–92%	BTS [15]	≥95%
6	Chest X-ray within 4 hours	NICE QS110 [5]	≥90%
7	Pulmonary rehab referral	BTS [6]	≥85%
8	Smoking cessation advice for smokers	USPSTF [16]	≥90%

NICE, National Institute of Health and Care Excellence; BTS, British Thoracic Society; GOLD, Global Initiative for Chronic Obstructive Lung Disease; USPSTF, United States Preventive Services Task Force. MDI = metered-dose inhaler. MDI with spacer was the preferred bronchodilator delivery method according to local orders; nebulized bronchodilator therapy was permitted only when MDI/spacer was contraindicated or not tolerated (e.g., inability to coordinate inhalation or severe agitation). Antibiotic decisions were made by treating physicians, guided by the order set criteria (see Methods), and were reviewed by the antimicrobial stewardship pharmacist.

from July to October 2024, followed by the identification of care gaps, the co-design of targeted interventions, and the implementation phase. The cycle concludes with the re-audit from January to March 2025 and subsequent planning for sustainability and scale-up. The figure illustrates how each phase informed the next, in line with the iterative principles of quality improvement methodology.

2.10. Measures

Eight process indicators were selected in accordance with international guidelines. These included: ABG within one hour of admission; nebulized bronchodilator administration within one hour; systemic corticosteroid prescription within four hours; appropriate antibiotic use based on clinical indication; oxygen therapy titrated to target saturation of 88–92% with documentation of FiO₂; chest X-ray within four hours; pulmonary rehab referral at discharge; and smoking cessation counseling for current smokers as shown in Table 1. Their guideline origins supported the validity of these measures, and data were abstracted using a structured form. Completeness and accuracy were assured through dual validation by two clinicians. Implementation fidelity and monitoring. We tracked fidelity using multiple process indicators: staff education attendance (percentage of targeted staff attending training sessions; 100% coverage reported); EHR log data confirming the display of the bedside prompt for triggered admissions; and order set opening/acceptance rates (EHR log metric). In 88% of triggered admissions, the order set was opened and used. These fidelity metrics are summarised in Supplementary Table S1.

2.11. Analysis

Descriptive statistics were used to summarize performance against each indicator. Pre- and post-intervention comparisons were made using percentage point differences. Confidence intervals were calculated for proportions. Given the sample size of 50 per audit cycle, we prioritized changes of more than 5% as potentially meaningful. Variations over time were examined through comparative tabulation. Analyses were primarily descriptive. For each process measure, we calculated proportions with 95% Wilson confidence intervals. For pre- and post-comparisons, we used two-sided Fisher's exact tests, given the sample sizes and the presence of small cell counts in some measures. We did not apply multiplicity correction because these analyses were exploratory and hypothesis-generating; we therefore interpret p-values cautiously, mindful of the potential for type I error across multiple comparisons. No formal prospective sample-size calculation was performed; this

activity was planned and conducted as a pilot/service evaluation to inform larger, more definitive future studies.

2.12. Ethical Considerations

The project was approved as a service evaluation and registered with the hospital's clinical audit department (Registration 2025-202). Formal ethics review was waived in accordance with institutional policy, as the project involved routine care data with no patient identifiers. The authors declared no conflicts of interest.

2.13. Reporting Standards

This quality improvement project is reported in accordance with the Standards for Quality Improvement Reporting Excellence (SQUIRE 2.0) guidelines [17]. The SQUIRE framework is specifically designed to guide the reporting of systematic efforts to improve healthcare quality, safety, and value in real-world clinical settings. A completed SQUIRE Checklist is provided in the supplementary materials to accompany this manuscript.

3. Results

3.1. Process Outcomes

At baseline (as shown in Table 2), compliance was highest for early administration of bronchodilators (92%) and initiation of corticosteroids (94%), indicating good adherence to acute pharmacotherapy. ABG sampling within one hour was achieved in 82% of patients. However, oxygen titration to SpO₂ 88–92% was documented in only 80% of cases, and only 64% of eligible patients were referred to pulmonary rehabilitation at discharge. The order set recommendation for a 5-day short course was followed in the majority of patients: 94% of treated patients received the recommended prednisolone 40 mg daily for 5 days. During the audit periods, 24 patients required noninvasive ventilation. A chest X-ray was obtained within four hours in 78% of cases. Among current smokers, documentation of cessation advice was noted in 87.5% of cases.

Following the interventions, compliance improved across all measures (shown in Table 3). The pulmonary rehab referral rate increased to 86%, meeting the target of 85% or higher. Oxygen titration documentation improved to 92%, reflecting gains but remaining just below the 95% goal. ABG and corticosteroid benchmarks were met, reaching 90% and 98%, respectively. Smoking cessation advice documentation exceeded the 90% target, rising to 95%.

Table 2: Baseline Audit Results (July–October 2024)

Standard	Target	n/N	Compliance % (95% CI)	Δ from Target
ABG within 1 hr.	≥90%	41/50	82.0% (69.0–90.6)	–8%
Nebulized bronchodilators <1 hr.	≥90%	46/50	92.0% (80.8–97.7)	+2%
Corticosteroids <4 hr.	≥95%	47/50	94.0% (83.5–98.8)	–1%
Appropriate antibiotics	≥90%	43/47	91.5% (79.6–97.6)	+1.5%
Oxygen titrated, SpO ₂ 88–92%	≥95%	40/50	80.0% (66.3–89.9)	–15%
Chest X-ray within 4 hr.	≥90%	39/50	78.0% (64.0–88.5)	–12%
Pulmonary rehab referral	≥85%	32/50	64.0% (49.2–77.1)	–21%
Smoking cessation advice	≥90%	35/40	87.5% (73.2–95.8)	–2.5%

Δ, change or delta; ABG, arterial blood gas; CI, confidence interval; hr, hour; n/N, number achieving target/total number assessed; SpO₂, peripheral capillary oxygen saturation.

Table 3: Re-Audit Results (January–March 2025)

Standard	Target	Baseline % (95% CI)	Re-audit % (95% CI)	Δ %	p value*	Target met?
ABG within 1 hr.	≥90%	82.0 (69.0–90.6)	90.0 (78.2–96.7)	+8.0	0.27	Yes
Nebs within 1 hr.	≥90%	92.0 (80.8–97.7)	94.0 (83.5–98.8)	+2.0	1.00	Yes
Corticosteroids within 4 hr.	≥95%	94.0 (83.5–98.8)	98.0 (89.4–99.9)	+4.0	0.61	Yes
Appropriate antibiotics	≥90%	91.5 (79.6–97.6)	94.0 (83.5–98.8)	+2.5	1.00	Yes
Oxygen titration	≥95%	80.0 (66.3–89.9)	92.0 (80.8–97.7)	+12.0	0.11	No
Chest X-ray within 4 hr.	≥90%	78.0 (64.0–88.5)	88.0 (75.7–95.5)	+10.0	0.28	No
Pulmonary rehab referral	≥85%	64.0 (49.2–77.1)	86.0 (73.3–94.2)	+22.0	0.020	Yes
Smoking cessation advice	≥90%	87.5 (73.2–95.8)	95.0 (83.1–99.4)	+7.5	0.43	Yes

Δ, change or delta; ABG, arterial blood gas; CI, confidence interval; hr, hour; Nebs, nebulized bronchodilators; p value, probability value.

To illustrate the comparative impact of our interventions across key clinical standards, a clustered bar chart (Figure 2) was constructed showing baseline and re-audit compliance percentages for all eight audit measures. This visualization highlights notable improvements in pulmonary rehabilitation referrals, corticosteroid timing, and smoking cessation documentation, with more modest gains in oxygen titration and imaging. While most standards exceeded their target thresholds in the re-audit cycle, a few still fell short, underscoring the importance of continued iterative improvements.

To summarize overall performance during the re-audit period, a pie chart (Figure 3) displays the proportion of care standards that met or did not meet predefined targets. Six of the eight standards achieved full compliance with benchmark criteria, demonstrating broad improvement across multiple domains of AECOPD care. The two remaining standards, oxygen titration and chest X-ray timing, showed progress but did not reach full compliance, indicating areas for further refinement.

To better understand the impact of each intervention and assess progress toward predefined goals, we tracked key performance indicators aligned with our action plan. These included oxygen titration compliance, staff education coverage, pulmonary rehabilitation referral rates, and smoking cessation documentation. For each domain, we compared baseline performance with targets and re-audit results. Table 4 summarizes these metrics and highlights which objectives were fully achieved and which remained areas for continued improvement.

The following bar chart (Figure 4) displays the trajectory of four key intervention metrics: baseline status, predefined targets, and post-intervention compliance. Improvements in staff education, smoking cessation documentation, and pulmonary rehab referral

rates all met or exceeded targets. Oxygen titration, while improved from baseline, fell slightly short of its target, highlighting an ongoing challenge in embedding oxygen therapy best practices into real-time clinical workflows.

3.2. Associations and Contextual Influences

The automated EHR prompts were consistently identified by staff as helpful in remembering care tasks during busy admissions. Feedback collected during clinical huddles confirmed that the oxygen checklist and discharge reminders resulted in improved documentation and care planning. Delays in chest X-ray performance were attributed to persistent bottlenecks in radiology transport and technician availability during peak hours. There were no unintended consequences, although two staff members noted occasional “alert fatigue” from excessive on-screen prompts.

4. Discussion

This quality improvement (QI) initiative demonstrated that structured, guideline-aligned interventions can meaningfully enhance the inpatient management of acute exacerbations of chronic obstructive pulmonary disease (AECOPD). By focusing on eight evidence-based process measures and implementing targeted changes across digital workflows, clinical education, and discharge planning, we were able to close care gaps within a short timeframe. The overall improvement observed in six of eight care standards underscores the feasibility and impact of low-cost, system-level interventions in improving care quality and equity for patients hospitalized with AECOPD.

One of the most striking findings was the substantial improvement in pulmonary rehabilitation (PR) referrals, which increased from

Table 4: Action Plan Metrics

Action Item	Baseline % (95% CI)	Target	Re-audit % (95% CI)	p value*	Status
Oxygen titration compliance	80.0 (66.3–89.9)	≥95%	92.0 (80.8–97.7)	0.11	Partial
Staff education coverage	0.0 (0.0–7.1)	100%	100.0 (92.9–100.0)	<0.001	Achieved
Pulmonary rehab referral rate	64.0 (49.2–77.1)	≥85%	86.0 (73.3–94.2)	0.020	Achieved
Smoking cessation documentation	87.5 (73.2–95.8)	≥95%	95.0 (83.1–99.4)	0.43	Achieved

CI, confidence interval; p value, probability value.

64% at baseline to 86% post-intervention, surpassing the 85% target. This success was directly attributable to the implementation of an automatic referral prompt within the electronic discharge summary. Historically, PR referrals are often neglected due to time constraints, lack of awareness, or assumptions about patient interest [18, 19, 20]. Our findings align with previous research showing that electronic reminders embedded within discharge workflows can significantly increase compliance with secondary prevention strategies in chronic disease care [21]. This is particularly important in COPD, where PR is strongly associated with reduced readmissions, improved exercise tolerance, and better quality of life [22, 23]. Importantly, the intervention helped eliminate variability based on individual clinician practices and supported more equitable care delivery across all patients.

Improvement in oxygen titration practices was another major focus of our initiative. At baseline, only 80% of patients received oxygen titrated to a target SpO₂ of 88–92% with documented FiO₂, far below the BTS-recommended 95% threshold [15]. Following the introduction of an oxygen prescription checklist in the EHR and targeted staff education, compliance improved to 92%. Although this fell just short of our goal, the 12-percentage point gain is clinically meaningful and represents a major step toward safer oxygen prescribing. The emphasis on titrated oxygen is clinically important. Studies [24,25] indicate that titrated oxygen therapy (target SpO₂ 88–92%) reduces the risk of hypercapnic respiratory failure and associated adverse outcomes compared with liberal high-flow oxygen in patients at risk of CO retention. Achieving and sustaining high compliance with oxygen targets therefore has direct safety implications for patients with AECOPD and remains a priority for future PDSA cycles [24, 25]. Our intervention directly addressed the root cause, lack of FiO₂ documentation and unclear targets, by prompting prescribers at the point of care. However, achieving full compliance will likely require continued reinforcement and monitoring, particularly during handoffs and night shifts when documentation lapses may occur.

The timing of core pharmacologic interventions also improved. Administration of systemic corticosteroids within four hours of admission rose from 94% to 98%, and timely initiation of bronchodilators remained above 90% across both audit cycles. These findings indicate a strong baseline awareness of pharmacologic protocols among frontline providers, which was further supported by our educational sessions. The importance of timely corticosteroids in reducing exacerbation duration and hospitalization length has been well described, and our near-universal compliance is encouraging [26, 27]. Similarly, appropriate antibiotic prescribing improved slightly, from 91.5% to 94%, reinforcing that clinicians are increasingly adopting an individualized approach based on purulent sputum and clinical infection markers, as recommended by GOLD and NICE.

Despite clear progress in several domains, two standards remained below target: chest X-ray completion within four hours (88% vs. 90% target) and oxygen titration compliance (92% vs. 95% target). The delay in chest imaging appears to be largely logistical. Informal feedback from nursing and medical staff pointed to porter availability and prioritization of other acutely unwell patients as contributing factors. Although we streamlined request pathways during the re-audit cycle, further improvements may require departmental-level coordination with radiology services. Introducing fast-track imaging protocols for medical admissions with respiratory compromise may be an effective next step.

The integration of a smoking cessation reminder into the discharge workflow proved highly successful, with documentation rates improving from 87.5% to 95%. This is consistent with prior work suggesting that digital nudges and automated prompts can reliably increase adherence to preventive counseling [28, 29]. Smoking remains the most significant modifiable risk factor in COPD progression, and inpatient encounters represent a critical opportunity for intervention [30, 31]. Our findings suggest that embedding such prompts into routine documentation processes minimizes omission and ensures consistency regardless of provider seniority or time constraints.

This project also contributes to the growing literature on digital quality improvement tools in internal medicine. Several studies have explored the impact of computerized decision support in chronic disease management, but relatively few have focused on acute care settings like AECOPD admissions [32, 33, 34, 35, 36, 37, 38, 39, 40]. Our experience highlights that with modest investment and stakeholder engagement, existing electronic health record platforms can be leveraged to embed real-time clinical decision support, improving both care delivery and equity. The interventions we implemented were designed to reduce variability stemming from individual clinician knowledge, shift patterns, or cognitive load—factors that disproportionately affect patients during unscheduled admissions. By standardizing care processes, we aim to reduce disparities and ensure that all patients receive evidence-based, guideline-concordant treatment.

5. Limitations

Several limitations warrant careful attention. First, this was a single-centre pilot before-and-after study with modest sample sizes ($n = 50$ per cycle); the study was not prospectively powered to detect small differences and may be at risk of type II error for some measures. Second, there was no randomization or concurrent control, so causal attribution to the interventions is limited; results may be affected by temporal confounding (seasonality, staffing patterns) or regression to the mean. Third, staff awareness of the project (Hawthorne effect) may have contributed to observed changes independent of the interventions. Fourth, although we

tracked several fidelity metrics, consistent application of the interventions across all shifts could not be guaranteed; orderset open/use rates are presented in Supplementary Table S1 and remain an area for ongoing monitoring. Fifth, the re-audit was performed shortly after implementation and may not reflect long-term sustainability; longer follow-up is planned. Finally, we performed multiple exploratory comparisons and did not adjust for multiplicity; readers should interpret p-values alongside effect sizes and confidence intervals. Although our initial intervention prioritised acute in-hospital process measures, we collected data on several secondary preventive items (vaccination status, smoking cessation counselling and documented inhaler checks) and identified remaining opportunities. The orderset already includes optional fields to facilitate vaccination checks, structured inhaler technique education (nursing checklist and leaflet), smoking cessation referral and automatic pulmonary rehabilitation referral; these items are pre-specified for inclusion in subsequent PDSA cycles. Future work will evaluate uptake, barriers to completion (for example time pressures and resource constraints), and impact on patient-centred outcomes such as readmissions and functional recovery.

Despite these limitations, the sustainability of our interventions is promising. Because the core changes were embedded in digital workflows and did not require additional staffing or new infrastructure, they are likely to persist over time with minimal maintenance. The success of the pulmonary rehab auto-referral and smoking cessation documentation prompts provides a clear model for replicating this approach in other chronic disease pathways, such as heart failure or diabetes care. Additionally, the strong staff engagement we observed—reflected in positive feedback during team huddles—suggests a fertile environment for ongoing quality improvement efforts.

Our QI project demonstrates that meaningful improvements in the quality, safety, and equity of AECOPD inpatient care can be achieved through pragmatic, system-level interventions. The gains in PR referral, oxygen safety, and preventive counseling highlight the power of digital tools to close longstanding care gaps. Although this project focused on process measures, many of the improvements we observed are strongly linked to favourable patient-centred outcomes in the literature. For example, pulmonary rehabilitation referrals are associated with reduced readmissions and improved exercise capacity and quality of life, and systematic smoking cessation counselling increases long-term abstinence and reduces disease progression [9-11, 30]. We therefore anticipate that improved uptake of these processes will translate into meaningful patient benefits; as such, subsequent PDSA cycles will include prospective measurement of patient-centred outcomes (for example 30- and 90-day readmission, length of stay, validated symptom scores and patient experience metrics) to quantify clinical impact beyond process adherence. Continued work is needed to optimize imaging workflows and further enhance oxygen prescribing practices. More broadly, our approach offers a replicable template for other internal medicine services seeking to bridge the gap between evidence and practice.

6. Conclusions

This QI project highlights how focused, system-integrated interventions can enhance the timeliness, quality, and equity of AE-COPD care. The work is likely to be sustainable given its reliance on EHR tools rather than high-resource staffing. With minimal adaptation, the approach could be extended to other respiratory or acute care pathways. Future studies should evaluate patient-centered outcomes such as readmission, recovery, and experience.

Continued refinement of digital tools and expansion to additional care settings are key next steps.

Conflicts of Interest

All authors declare that they have no conflicts of interest, financial or otherwise, relevant to the content of this manuscript. The authors affirm that they have no affiliations with or involvement in any organization or entity with any financial interest or non-financial interest in the subject matter discussed in this manuscript.

Funding Source

This project did not receive any external funding. All work was conducted as part of internal departmental quality improvement activities, supported by existing institutional resources. The funding organization (i.e., the hospital or department) had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgments

We thank the Clinical Informatics team, Pulmonary Rehabilitation Unit, and General Medicine nursing staff for their support in implementing the project interventions. We are especially grateful to the Clinical Audit Department for providing oversight, audit templates, and governance support. Their contributions were essential to the successful completion of this quality improvement initiative.

Institutional Review Board (IRB)

This study was reviewed by the Clinical Audit and Quality Governance Department at a tertiary care hospital (Registration 2025-202) and approved as a quality improvement initiative. In accordance with national and institutional policies, it did not require approval from a Research Ethics Committee or Institutional Review Board, as it involved retrospective and prospective review of anonymized data from routine clinical care. Written informed consent was not required as per institutional policy for service evaluation projects that do not involve deviation from standard care.

Large Language Model

None

Authors Contribution

All authors contributed substantially and equally to the conception, design, execution, and reporting of this quality improvement project. KA, AA, and MS led the development of the project protocol, coordinated data collection, and facilitated intervention implementation. MI and HS contributed to critical data analysis, interpretation of findings, and literature review. MS drafted the initial manuscript with input from all co-authors. All authors participated in revising the manuscript for important intellectual content, approved the final version for submission, and agree to be accountable for all aspects of the work to ensure accuracy and integrity. No professional medical writers or artificial intelligence generative processing methods were involved. All authors meet the International Committee of Medical Journal Editors (ICMJE) authorship criteria and adhere to the ethical standards outlined by the Committee on Publication Ethics (COPE).

Data Availability

The data that support the findings of this study are available from the corresponding author upon reasonable request. De-identified patient-level data used in the audit and quality improvement analysis are retained in secure institutional repositories and may be shared with qualified researchers, subject to institutional approvals and data-sharing agreements. No individual-level patient-identifiable information is publicly available in compliance with institutional and ethical policies.

References

1. Chow L, Parulekar AD, Hanania NA. Hospital management of acute exacerbations of chronic obstructive pulmonary disease. *J Hosp Med*. 2015;10(5):328-39. [PMID: 25820201, <https://doi.org/10.1002/jhm.2334>].
2. Chow R, So OW, Im JHB, Chapman KR, Orchanian-Cheff A, Gershon AS, et al. Predictors of Readmission, for Patients with Chronic Obstructive Pulmonary Disease (COPD) - A Systematic Review. *Int J Chron Obstruct Pulmon Dis*. 2023;18:2581-617. [PMID: 38022828, PMCID: PMC10664718, <https://doi.org/10.2147/COPD.S418295>].
3. Langfeld K. World Lung Day 2019: Respiratory Groups Unite to Call for Healthy Lungs for All; 2019.
4. Press VG, Au DH, Bourbeau J, Dransfield MT, Gershon AS, Krishnan JA, et al. Reducing Chronic Obstructive Pulmonary Disease Hospital Readmissions. An Official American Thoracic Society Workshop Report. *Ann Am Thorac Soc*. 2019;16(2):161-70. [PMID: 30707066, PMCID: PMC6812156, <https://doi.org/10.1513/AnnalsATS.201811-755WS>].
5. QS110 Pneumonia in adults. National Institute of Health and Care Excellence;.
6. Bolton CE, Bevan-Smith EF, Blakey JD, Crowe P, Elkin SL, Garrod R, et al. British Thoracic Society guideline on pulmonary rehabilitation in adults: accredited by NICE. *Thorax*. 2013;68(Suppl 2):ii1-ii30. [<https://doi.org/10.1136/thoraxjnl-2013-203808>].
7. National Institute for Health and Care Excellence. NG115 Chronic obstructive pulmonary disease in over 16s: diagnosis and management. National Institute for Health and Care Excellence; 2019.
8. Kostikas K, Hillas G, Gogali A. 2025 GOLD Report: What is New and What is Noteworthy for the Practicing Clinician. *COPD*. 2025;22(1):2451613. [PMID: 39912681, <https://doi.org/10.1080/15412555.2025.2451613>].
9. Aldhahir AM, Alghamdi SM, Alqahtani JS, Alqahtani KA, Al Rajah AM, Alkathlan BS, et al. Pulmonary rehabilitation for COPD: A narrative review and call for further implementation in Saudi Arabia. *Ann Thorac Med*. 2021;16(4):299-305. [PMID: 34820017, PMCID: PMC8588944, https://doi.org/10.4103/atm.atm_639_20].
10. Cox NS, Oliveira CC, Lahham A, Holland AE. Pulmonary rehabilitation referral and participation are commonly influenced by environment, knowledge, and beliefs about consequences: a systematic review using the Theoretical Domains Framework. *J Physiother*. 2017;63(2):84-93. [PMID: 28433238, <https://doi.org/10.1016/j.jphys.2017.02.002>].
11. Lusuardi M, Garuti G, Massobrio M, Spagnolatti L, Borciani N. The audit-intervention-reaudit cycle can improve diagnostic standards for chronic obstructive pulmonary disease (COPD). *Intern Emerg Med*. 2014;9(1):113-4. [PMID: 23636565, <https://doi.org/10.1007/s11739-013-0952-8>].
12. Man WD, Puhan MA, Harrison SL, Jordan RE, Quint JK, Singh SJ. Pulmonary rehabilitation and severe exacerbations of COPD: solution or white elephant? *ERJ Open Res*. 2015;1(2):00050-2015. [PMID: 27730157, PMCID: PMC5005122, <https://doi.org/10.1183/23120541.00050-2015>].
13. Price LC, Lowe D, Hosker HS, Anstey K, Pearson MG, Roberts CM, et al. UK National COPD Audit 2003: Impact of hospital resources and organisation of care on patient outcome following admission for acute COPD exacerbation. *Thorax*. 2006;61(10):837-42. [PMID: 16449268, PMCID: PMC2104768, <https://doi.org/10.1136/thx.2005.049940>].
14. Suarez-Gonzalez P, Suarez-Elosegui A, Arias-Fernandez L, Perez-Regueiro I, Jimeno-Demuth FJ, Lana A. Nursing diagnoses and hospital readmission of patients with respiratory diseases: Findings from a case-control study. *Nurs Open*. 2024;11(5):e2182. [PMID: 38783599, PMCID: PMC11116758, <https://doi.org/10.1002/nop.2.2182>].
15. O'Driscoll BR, Howard LS, Earis J, Mak V, British Thoracic Society Emergency Oxygen Guideline G, Group BTSEOGD. BTS guideline for oxygen use in adults in healthcare and emergency settings. *Thorax*. 2017;72(Suppl 1):ii1-ii90. [PMID: 28507176, <https://doi.org/10.1136/thoraxjnl-2016-209729>].
16. US Preventive Services Task Force. Interventions for Tobacco Smoking Cessation in Adults, Including Pregnant Persons: US Preventive Services Task Force Recommendation Statement. *JAMA*. 2021;325(3):265-79. [<https://doi.org/10.1001/jama.2020.25019>].
17. Ogrinc G, Davies L, Goodman D, Batalden P, Davidoff F, Stevens D. SQUIRE 2.0 (Standards for QUality Improvement Reporting Excellence): revised publication guidelines from a detailed consensus process. *BMJ Quality & Safety*. 2016;25(12):986-92. [<https://doi.org/10.1136/bmjqs-2015-004411>].
18. Augustine A, Bhat A, Vaishali K, Magazine R. Barriers to pulmonary rehabilitation - A narrative review and perspectives from a few stakeholders. *Lung India*. 2021;38(1):59-63. [PMID: 33402639, PMCID: PMC8066922, https://doi.org/10.4103/lungindia.lungindia_116_20].
19. Qi J, Cai Q, Zhang W, Gu Y. Effects of pulmonary telerehabilitation in patients with chronic obstructive pulmonary disease. *Chin Med J (Engl)*. 2025;138(12):1493-5. [PMID: 40413618, PMCID: PMC12180807, <https://doi.org/10.1097/CM9.0000000000003650>].
20. Thakor M, Singh V, Manifield J, Orme MW, Bhardwaj P, Chauhan NK, et al. Community-Based Pulmonary Rehabilitation in an Economically Deprived Area of Jodhpur, India: A Mixed-Methods Feasibility Trial. *Int J Chron Obstruct Pulmon Dis*. 2025;20:473-8. [PMID: 40041473, PMCID: PMC11878114, <https://doi.org/10.2147/COPD.S488766>].
21. Willis VC, Thomas Craig KJ, Jabbarpour Y, Scheufele EL, Arriaga YE, Ajinkya M, et al. Digital Health Interventions to Enhance Prevention in Primary Care: Scoping Review. *JMIR Med Inform*. 2022;10(1):e33518. [PMID: 35060909, PMCID: PMC8817213, <https://doi.org/10.2196/33518>].
22. Press VG, Nici L. Improving Pulmonary Rehabilitation Participation: It's Time to Examine the System. *Ann Am Thorac Soc*. 2024;21(12):1655-6. [PMID: 39601503, PMCID: PMC11622825, <https://doi.org/10.1513/AnnalsATS.202410-1003ED>].
23. Rochester CL, Alison JA, Carlin B, Jenkins AR, Cox NS, Bauldoff G, et al. Pulmonary Rehabilitation for Adults with Chronic Respiratory Disease: An Official American Thoracic Society Clinical Practice Guideline. *Am J Respir Crit Care Med*. 2023;208(4):e7-e26. [PMID: 37581410, PMCID: PMC10449064, <https://doi.org/10.1164/rccm.202306-1066ST>].
24. Echevarria C, Steer J, Wason J, Bourke S. Oxygen therapy and inpatient mortality in COPD exacerbation. *Emerg Med J*. 2021;38(3):170-7. [PMID: 33243839, <https://doi.org/10.1136/emered-2019-209257>].
25. Kelly CA, Maden M. How do health-care professionals perceive oxygen therapy? A critical interpretive synthesis of the literature. *Chron Respir Dis*. 2015;12(1):11-23. [PMID: 25492978, <https://doi.org/10.1177/1479972314562408>].
26. Putcha N, Wise RA. Medication Regimens for Managing COPD Exacerbations. *Respir Care*. 2018;63(6):773-82. [PMID: 29794210, PMCID: PMC6818681, <https://doi.org/10.4187/respcare.05912>].
27. Woods JA, Wheeler JS, Finch CK, Pinner NA. Corticosteroids in the treatment of acute exacerbations of chronic obstructive pulmonary disease. *Int J Chron Obstruct Pulmon Dis*. 2014;9:421-30. [PMID: 24833897, PMCID: PMC4014384, <https://doi.org/10.2147/COPD.S51012>].
28. Nurchis MC, Di Pumpo M, Perilli A, Greco G, Damiani G. Nudging Interventions on Alcohol and Tobacco Consumption in Adults: A Scoping Review of the Literature. *Int J Environ Res Public Health*. 2023;20(3). [PMID: 36767077, PMCID: PMC9913966, <https://doi.org/10.3390/ijerph20031675>].
29. Sumner J, Bundale A, Lim HW, Phan P, Motani M, Mukhopadhyay A. Developing an Artificial Intelligence-Driven Nudge Intervention

- to Improve Medication Adherence: A Human-Centred Design Approach. *J Med Syst.* 2023;48(1):3. [PMID: 38063940, PMCID: PMC10709244, <https://doi.org/10.1007/s10916-023-02024-0>].
30. Paulin LM, Halenar MJ, Edwards KC, Lauten K, Taylor K, Brunette M, et al. Relationship Between Tobacco Product Use and Health-Related Quality of Life Among Individuals With COPD in Waves 1-5 (2013-2019) of the Population Assessment of Tobacco and Health Study. *Chronic Obstr Pulm Dis.* 2024;11(1):68-82. [PMID: 38113525, PMCID: PMC10913919, <https://doi.org/10.15326/jcopdf.2023.0422>].
31. Wu J, Sin DD. Improved patient outcome with smoking cessation: when is it too late? *Int J Chron Obstruct Pulmon Dis.* 2011;6:259-67. [PMID: 21814462, PMCID: PMC3144846, <https://doi.org/10.2147/COPD.S10771>].
32. Chen W, Howard K, Gorham G, O'Bryan CM, Coffey P, Balasubramanya B, et al. Design, effectiveness, and economic outcomes of contemporary chronic disease clinical decision support systems: a systematic review and meta-analysis. *J Am Med Inform Assoc.* 2022;29(10):1757-72. [PMID: 35818299, PMCID: PMC9471723, <https://doi.org/10.1093/jamia/ocac110>].
33. Dijk SW, Wollny C, Barkhausen J, Jansen O, Mildenerger P, Halfmann MC, et al. Evaluation of a Clinical Decision Support System for Imaging Requests: A Cluster Randomized Clinical Trial. *JAMA.* 2025;333(14):1212-21. [PMID: 39928308, PMCID: PMC11811869, <https://doi.org/10.1001/jama.2024.27853>].
34. Fujita K, Hooper P, Masnoon N, Lo S, Gnjjidic D, Etherton-Beer C, et al. Impact of a Comprehensive Intervention Bundle Including the Drug Burden Index on Deprescribing Anticholinergic and Sedative Drugs in Older Acute Inpatients: A Non-randomised Controlled Before-and-After Pilot Study. *Drugs Aging.* 2023;40(7):633-42. [PMID: 37160561, PMCID: PMC10299923, <https://doi.org/10.1007/s40266-023-01032-6>].
35. Huang S, Liang Y, Li J, Li X. Applications of Clinical Decision Support Systems in Diabetes Care: Scoping Review. *J Med Internet Res.* 2023;25:e51024. [PMID: 38064249, PMCID: PMC10746969, <https://doi.org/10.2196/51024>].
36. Roshanov PS, Misra S, Gerstein HC, Garg AX, Sebaldt RJ, Mackay JA, et al. Computerized clinical decision support systems for chronic disease management: a decision-maker-researcher partnership systematic review. *Implement Sci.* 2011;6:92. [PMID: 21824386, PMCID: PMC3170626, <https://doi.org/10.1186/1748-5908-6-92>].
37. Sommers S, Tolle H, Napier C, Hoppe J. Targeted messaging to improve the adoption of clinical decision support for prescription drug monitoring program use. *J Am Med Inform Assoc.* 2023;30(10):1711-6. [PMID: 37433582, PMCID: PMC10531197, <https://doi.org/10.1093/jamia/ocad127>].
38. Velickovski F, Ceccaroni L, Roca J, Burgos F, Galdiz JB, Marina N, et al. Clinical Decision Support Systems (CDSS) for preventive management of COPD patients. *J Transl Med.* 2014;12 Suppl 2(Suppl 2):S9. [PMID: 25471545, PMCID: PMC4255917, <https://doi.org/10.1186/1479-5876-12-S2-S9>].
39. Ventura F, Sousa P, Dixe MA, Ferreira P, Martinho R, Dias SS, et al. A Clinical Decision Support System for Remote Monitoring of Cardiovascular Disease Patients: A Clinical Study Protocol. *Front Public Health.* 2022;10:859890. [PMID: 35615041, PMCID: PMC9124932, <https://doi.org/10.3389/fpubh.2022.859890>].
40. Xi H, Zhang Y, Zuo R, Li W, Zhang C, Sun Y, et al. Forecasting Hospitalization for Adult Asthma Patients in Emergency Departments Based on Multiple Environmental and Clinical Factors. *J Asthma Allergy.* 2025;18:861-76. [PMID: 40469908, PMCID: PMC12135953, <https://doi.org/10.2147/JAA.S512405>].